



ALLIANCE™

(<https://www.globalseafood.org>).



 Aquafeeds

Algae alternative: Chlorella studied as protein source in tilapia feeds

3 May 2013

By Ingrid Lupatsch, Ph.D. and Chris Blake

Fish achieved satisfactory growth, with FCR efficiency comparable to soybean-based feed



Experimental fish feeds containing different proportions of dried *Chlorella* in substitution for fishmeal.

In both aquaculture and agriculture, producers commonly rely on formulated feeds to ensure optimal growth, health and quality of the farmed animals. Fishmeal and fish oil from wild fisheries have traditionally been major constituents of aquafeeds, but their supply is finite. Therefore, raw ingredients other than fishmeal are being selected for their nutritive value, balance of amino acids, protein digestibility, quality of lipids and fatty acids, availability and cost.

Algae alternatives

Algae biomass is being considered as one of the alternative ingredients of the future. The composition of micro- and macroalgae vary considerably among species, but also depend upon culture conditions. The average protein level in macroalgae is around 8 to 15 percent dry matter, while the average lipid content is only 1 to 3 percent. This compares to a protein content of 30 to 50 percent dry matter for microalgae, which have lipid contents as high as 40 percent.

Freshwater algae such as *Chlorella* and *Spirulina* seem to have good potential as protein sources, whereas marine microalgae are the fundamental source of the long-chain polyunsaturated fatty acids that are crucial for human health, as well as that of aquaculture animals.

In a study, the authors examined the efficacy and nutritional properties of *Chlorella vulgaris* as an alternative feed ingredient and protein source for the culture of all-male Nile tilapia, *Oreochromis niloticus*.



(<https://events.globalseafood.org/responsible-seafood-summit>).

Experimental setup

Genetically male tilapia sourced as fry and raised at the Centre for Sustainable Aquaculture Research at Swansea University in the United Kingdom were used for all the trials. The trials were set up indoors as part of a freshwater recirculation system, which included mechanical and biofiltration units, a protein skimmer and a sand filter. Water temperature was kept at 27 degrees-C, and photoperiod was set at 12 hours of light daily. Temperature and dissolved-oxygen levels were measured daily, while total ammonia nitrogen, nitrite, nitrate and pH were measured weekly.

Feeds were prepared by mixing the dry ingredients with binder and water, and extruding the feed through a meat grinder, followed by thorough drying. The resulting pellets had a diameter of 2.5 mm and were stable up to 24 hours in water.

Digestibility trial

Digestibility studies of the algae biomass were performed by adding an indigestible marker (chromic oxide) to the feed and collecting the fecal matter by siphoning. By assessing the ratio of marker to energy or nutrient in the feed compared to their ratio in fecal matter, the digestibility of the feed could be established.

Fishmeal was used as the reference ingredient with the test diet mixed at 50 percent fishmeal and 50 percent *Chlorella*. Fifteen tilapia weighing on average 250 grams each were stocked per tank. Fecal matter from each tank was pooled over the trial period until sufficient fecal matter had been collected for analyses. The digestibility of the ingredients, calculated using well-established equations, is presented in Table 1.

Lupatsch, Apparent digestibility, Table 1

	Dry Matter (%)	Protein (%)	Organic Matter (%)	Energy (%)
Fishmeal*	73.6	90.5	82.1	83.4
Chlorella**	50.1	63.5	58.1	59.1

* Composition of fishmeal/kg as fed: dry matter 944 g, crude protein 645 g, lipid 99 g, ash 194 g, gross energy 18.82 MJ.

** Composition of Chlorella/kg as fed: dry matter 983 g, crude protein 472 g, lipid 82 g, ash 82 g, gross energy 21.14 MJ.

Table 1. Apparent digestibility coefficients of Chlorella and fishmeal in tilapia.

Growth trial

Fifteen tilapia of 35 g initial size were stocked in 150-L tanks. Feeds were formulated to contain 40 percent protein and 9 percent lipid, and to gradually include *Chlorella* at the expense of fishmeal (Table 2). Fish were fed manually to apparent satiation up to four times daily. Any uneaten pellets were collected at the end of the day.

Lupatsch, Formulation (g/kg as fed), Table 2

Ingredient	Fishmeal	30% <i>Chlorella</i>	60% <i>Chlorella</i>	100% <i>Chlorella</i>
Formulation				
Fishmeal	640	420	210	–
<i>Chlorella</i>	–	260	520	780
Corn starch	300	250	180	120
Vitamins, minerals	10	10	10	10
Dicalcium phosphate	–	10	30	50
Vegetable oil	20	20	20	20
Binder	30	30	30	20
Composition				
Dry matter (g)	930	920	930	930

Ash (g)	119	101	95	96
Lipid (g)	84	85	87	88
Crude protein (g)	413	394	382	377
Gross energy (MJ)	18.01	18.19	18.10	19.15
Digestible protein* (g)	374	323	279	239
Digestible energy* (MJ)	14.86	13.70	12.46	12.00
Digestible protein/Digestible energy ratio (g/MJ)	25.1	23.6	22.4	19.9

* Incorporating results from digestibility trial.

Table 2. Formulation (g/kg as fed) and composition of experimental feeds.

Feed intake was quantified and evaluated in relation to growth response. Fish were sampled initially and at the end of the 31-day growth trial for further chemical analyses. Through comparative body composition of fish carcass, the relationships between dietary protein and energy intake, and protein and energy deposition were assessed, which allowed estimation of the utilization efficiency of *Chlorella*.

Results

Results indicated that feed efficiency and growth performance were best for the fishmeal control and gradually deteriorated with increasing inclusion levels of *Chlorella* (Table 3 and Fig. 1). The feeds containing *Chlorella* were well accepted by tilapia, and feed intake was initially increased to make up for the low contents of digestible protein and energy (Table 3). However, the tilapia were apparently reaching the limit of how much feed they could consume.

Lupatsch, Performance parameters, Table 3

Treatment	Initial Weight (g/fish)	Final Weight (g/fish)	Specific Growth Rate	Feed Intake (%/day)	Feed-Conversion Ratio
Fishmeal	35.3	101.6	3.40	3.56	1.00
30% <i>Chlorella</i>	35.2	97.5	3.29	4.50	1.31
60% <i>Chlorella</i>	35.3	89.9	3.00	4.42	1.44
100% <i>Chlorella</i>	36.0	84.5	2.75	4.49	1.58

Table 3. Performance parameters of tilapia after 31 days of growth at 27 degrees C.

The daily feed intake and daily weight gain of fish fed the experimental diets are illustrated in Fig. 1. Despite the higher feed intake for the feeds with algae, daily weight gain decreased with increasing dietary inclusion of *Chlorella*. The result was rising feed conversion, meaning more feed was needed to

produce one unit of weight gain.

Fig. 1: Relationship of feed intake to resulting weight gain in tilapia.

Perspectives

Chlorella vulgaris shows some potential as an alternative feed ingredient in aquaculture. Although the algal feeds were not utilized as efficiently as the fishmeal feed, *Chlorella* was well accepted. Overall, tilapia were able to achieve satisfactory growth on the feed with algae, and feed-conversion efficiency was comparable to that for a soybean-based feed.

A possible means of increasing the nutritional value of algal biomass would be to break down the cell wall fragments by mechanical treatment or even removal of most of the fiber, although such additional processing steps may be too expensive. At present, the costs of fishmeal and fish oil are steadily increasing. However, all categories of algal products are currently much higher in cost than the commodity feedstuffs used in aquafeeds.

(Editor's Note: This article was originally published in the May/June 2013 print edition of the Global Aquaculture Advocate.)

Authors



INGRID LUPATSCH, PH.D.

Center for Sustainable Aquaculture Research
Swansea University
Singleton Park, Swansea SA2 8PP, United Kingdom

i.lupatsch@swansea.ac.uk (<mailto:i.lupatsch@swansea.ac.uk>)



CHRIS BLAKE

Postgraduate Student
Center for Sustainable Aquaculture Research
Swansea University
Singleton Park, Swansea SA2 8PP, United Kingdom

Copyright © 2023 Global Seafood Alliance

All rights reserved.