



ALLIANCE™

(<https://www.globalseafood.org>).



Health &
Welfare

Aquaculture and *Aeromonas hydrophila*: A complex interplay of environmental factors and virulence

15 July 2024

By Darryl Jory, Ph.D.

Study reviews the pathogenic bacterium *A. hydrophila*, its impact on aquacultured fish and its relationship with environmental factors



Aquaculture faces significant challenges due to bacterial pathogens like *Aeromonas hydrophila*, which can severely impact production and fish health. Understanding the relationship between environmental factors, host susceptibility, and bacterial virulence is crucial for effectively managing and mitigating the risks associated with *A. hydrophila* in aquaculture systems. Improved knowledge of its virulence and interplay with environmental factors paves the way for developing strategies to prevent and control diseases in aquaculture. Photo of *A. hydrophila* colonies (growing on the blood agar; colonies shown with reflected light) by Stefan Walkowski (CC BY-SA 4.0 <https://creativecommons.org/licenses/by-sa/4.0>, via Wikimedia Commons).

A **recent study** (<https://doi.org/10.1007/s10499-024-01535-y>) conducted by Dr. Bahaa Abdella and co-workers at Kafrelsheikh University in Egypt and published in *Aquaculture International* discusses the pathogenic bacterium *Aeromonas hydrophila*, its impact on aquacultured fish and its relationship with various environmental factors.

“Our research highlights the importance of understanding how environmental conditions influence the virulence of *Aeromonas hydrophila* in aquaculture. It involves a complex and dynamic interaction between the bacterium and its environment. Key points provided include monitoring temperature, pH

levels, and ammonia concentrations to prevent disease outbreaks. Proper management of these factors can improve fish health and reduce bacterial infections” stated to the *Advocate* Dr. Bahaa Abdella from Kafrelsheikh University in Egypt and senior author of the study.

Among **diseases** (<https://doi.org/10.1292/jvms.19-0025>), caused by *Aeromonas* spp. in fish are hemorrhagic septicemia, ulcer disease, and motile *Aeromonas* septicemia. These diseases can result in **high mortality rates and high economic losses** (https://doi.org/10.4194/2459-1831-v3_1_03) in aquaculture. *A. hydrophilic* is the predominant disease-causing species in this **bacterial genus** (<https://doi.org/10.1016/j.micpath.2020.104424>), in cultured freshwater fish and other species on a global scale. Its genetic diversity makes it difficult to serotype or genotype the strains of *A. hydrophila* due to the high complexity in terms of the type and the number of **virulence genes present** (<https://doi.org/10.1038/s41598-023-34887-1>) in each isolate. Besides being a fish pathogen, *A. hydrophila* can also **cause diseases in humans** (<https://doi.org/10.1128/CMR.00039-09>) – such as gastroenteritis, wound infections, and septicemia in immunocompromised individuals.

Environmental factors such as mineral availability, salinity, dissolved oxygen concentration, pH value and temperature – as well as poor management (malnutrition, overfeeding and overcrowding) – in hatchery facilities may cause stress to cultured animals and make them **more susceptible to infections** (<https://doi.org/10.1186/s12866-023-02827-8>) by *A. hydrophila*.



(<https://chickenofthesea.com/shrimpwithapurpose/>).

Moreover, *A. hydrophila* genomes are encoding different types of virulence factors. The expression and regulation of these genes may vary depending on the bacterial strain, host, and environmental conditions. It is not fully understood how environmental conditions have an effect on intrinsic bacterial characteristics and the **expression of virulence genes** (<https://doi.org/10.2478/s11756-020-00609-5>), so it is an open question. Thus, understanding the diversity, function, and environmental regulation of these genes is crucial for developing effective strategies to prevent and treat *A. hydrophila* infections in fish farms.

“*A. hydrophila* poses significant challenges to aquaculture because of its ability to thrive in diverse environmental conditions and infect a wide range of hosts. Factors such as temperature, dissolved oxygen, pH, ammonia, and nutrient levels influence the expression of virulence genes and toxin production. Understanding these mechanisms is critical to developing strategies to control aquatic bacterial infections and promote healthier and more sustainable aquaculture practices,” according to Dr. Abdella.



Nanoparticles are a promising alternative for the management and mitigation of vibriosis in aquaculture

Nanoparticles may have potential as antibacterial drugs for Vibriosis control through their various mechanisms of action against bacteria.



Global Seafood Alliance

Physical factors influence the microbial growth rate, such as temperature and pH value. However, little is known about how these factors can modify the expression of the virulence genes in the pathogenic microbes. It is critical to understand what triggers virulence factor expression or at least the most essential one in a specific pathogen.

For example, temperature-dependent expression of virulence genes in *Aeromonas* spp. in fish has been reported for some genes. However, the molecular mechanisms and regulatory pathways involved in this process are not fully understood. Adding to that, the increase in temperature is found to be stressful to the host, which increases the expression of certain steroids which increase the vulnerability of the fish for infection.

Fluctuations in temperature – and also pH – significantly influence the metabolic activity and growth rate of *A. hydrophila*, thereby modulating its virulence and overall pathogenicity. Overall, the **expression of virulence genes** (<https://doi.org/10.3389/fpubh.2021.692166>) in Aeromonads can vary depending on the water temperature and the host animal, and the inconsistency of the expression patterns of many virulence factors in the same strain in response to temperature changes emphasizes the difficulty and the need to identify the major virulence factor that is responsible for disease development in the susceptible host.

Aeromonads are facultative anaerobes, and as all other facultative microbes, they may switch between aerobic and anaerobic metabolism. Such switching behaviors are controlled by gene expression and are triggered by molecular sensing of oxygen concentration. No surprise, it has developed a molecular mechanism for doing so through gene expression control, including virulence genes. Oxygen can regulate virulence genes by modulating the activity of transcription factors, which are proteins that bind to specific DNA sequences and control gene expression.

Overall, oxygen may be considered as a key environmental factor that influences the expression of virulence genes in many bacterial strains including *A. hydrophila* and their ability to cause disease. However, there has been little research into the effect of oxygen concentrations on the expression of virulence genes in fish infections. Fig. 1 depicts the interplay between the environment, microbes and the host.




Fig. 1: The interactive relationship between environmental factors, microbes and the host.

The **formulation of fish diets** (<https://doi.org/10.1038/srep35232>) has a considerable impact on microbial ecology in the aquatic environment, as well as microbial pathogenicity, interaction and abundance, and *A. hydrophila* infection is associated with water with a high organic load and nutrient-rich feed pollutes aquatic water systems. The unutilized diet contributes to an increase in aeromonads in the pond water.

The precise formulation of the fish feed and its utilization by fish is very critical in controlling the infection and suppressing the activation of virulence genes in *A. hydrophila*. More research is needed to confirm the link between various dietary components and virulence gene overexpression in *A. hydrophila*. However, feeding fish a diet high in omega-3 fatty acids can help improve the immune system and make the fish less susceptible to infection; this relationship is not fully understood.

The environmental regulation of virulence in *A. hydrophila* emerges as a complex and dynamic interplay between the bacterium and its environment. Temperature, dissolved oxygen, pH, ammonia and nutrient availability act as modulators, coordinating the expression of virulence genes and toxin production. This remarkable adaptability allows *A. hydrophila* to thrive in diverse aquatic environments and establish infections across a range of hosts, posing a significant challenge to aquaculture.

“Unraveling the mechanisms of environmental virulence regulation presents a compelling target for future research. By elucidating these pathways, we can pave the way for the development of targeted interventions to mitigate *A. hydrophila* infections, ultimately promoting a healthier and more sustainable aquaculture industry,” concluded the study authors.

[Read the full study.](https://doi.org/10.1007/s10499-024-01535-y) (<https://doi.org/10.1007/s10499-024-01535-y>).

[@GSA_Advocate](https://twitter.com/GSA_Advocate) (https://twitter.com/GSA_Advocate).

Author



DARRYL JORY, PH.D.

Editor Emeritus
Responsible Seafood Advocate

darryl.jory@globalseafood.org (<mailto:darryl.jory@globalseafood.org>).

Copyright © 2025 Global Seafood Alliance

All rights reserved.