



[FEED SUSTAINABILITY \(/ADVOCATE/CATEGORY/FEED-SUSTAINABILITY\)](#)

Maximizing nutrition for adult marine fish

Thursday, 12 November 2015

By **Jorge A. Suarez, Ph.D.** and **Carlos Tudela, Christina Belfranin, Steven Sutton, Matthew Taynor, Ronald Hoenig, Drew Davis, Thiago Raggi, Daniel D. Benetti, Ph.D.**

Digestible protein-digestible energy ratios key to optimizing feeding efficiency

Feeds can represent more than 70 percent of the production costs in aquaculture. Since increased consumption of feed occurs in the final stages of the growout cycle, it is clear the development of ecologically efficient, economically viable and nutritionally adequate diets for adult marine fish should be a priority.

Yet we are still faced with insufficient information on the nutrition requirements of large sizes of commercially important tropical marine fish species and no clear information on the maximum levels of replacement of fishmeal and fish oil in diets for near-market-size fish. Feed manufacturers have made important progress towards formulating and developing better diets, but additional information is required to improve the sustainability and economic viability of the industry.

The aquaculture nutrition program at the University of Miami in Florida, USA, was established in 2010 to provide scientific and technological support in the area of nutrition to companies dedicated to the aquaculture of tropical marine fish. With the support of the United Soybean Board, Illinois Soybean Association, U.S. Soybean Export Council and Open Blue Sea Farms in Panama, the program has conducted research on nutritional requirements and traits of feed ingredients for cobia, pompano and, with Martec S.A. in Costa Rica, spotted rose snappers.



Adult Florida pompano were weighed and measured during harvest.

Fishmeal replacement for adult fish

A study on the effects of replacing dietary fishmeal with soy-based products on the production performance of cobia between 1.8 and 3.2 kg was conducted by Suarez et al. (2013. *Aquaculture* 416-417. 328-333). The soy-based products used were dehulled soybean meal, solvent-extracted soybean meal and a novel non-genetically modified variety of soybean meal.

Results indicated that up to 80 percent of the fishmeal could be replaced in larger cobia, attaining a fish in:fish out ratio of 1.3 without compromising growth performance or health (Table 1). The ability to replace such a high amount of fishmeal indicated the nutritional requirements of cobia may change with age and that developing more cost-effective and environmentally sustainable diets is possible without compromising health or growth rates.

This study helped validate the need for additional research on cobia and others species, not just at the juvenile stage, but at every stage of the commercial production chain.

Dietary treatment	Final weight (kg)	Weight gain (%)	Specific growth rate (%/day)	Feed efficiency	Feed conversion ratio	Fish in:fish out
Days 1-29*						
1	2.38	35.5a	1.02a	0.56	1.80	2.08c
2	2.58	24.4b	0.74b	0.51	1.95	1.17b
3	2.50	35.6a	1.02a	0.57	1.81	1.10b
4	2.38	33.1a	0.97a	0.57	1.74	0.91a
Pr>F (spell out)	0.24	0.014	0.008	0.82	0.87	0.003
Pooled standard error	0.46	15.04	0.37	0.08	0.28	0.52
Days 2-60**						
1	2.80	56.2	0.76a	0.49	2.0	2.30c
2	3.00	49.9	0.61b	0.47	2.1	1.27b

Digestible protein-digestible energy ratios

In a study in review with adult Florida pompano by Matthew Taylor, the effects of dietary digestible protein:digestible energy (D.P.:D.E.) ratios on growth, feed efficiency, nitrogen utilization and body composition were investigated. A growth trial and subsequent digestibility trial were carried out utilizing five diets based on anchovy meal, soybean meal or corn gluten meal as protein sources. The feeds were formulated to have different D.P.:D.E. ratios ranging 21.7-26.3 mg protein/kJ.

Fish were fed to apparent satiation by hand once daily at 9 a.m. for 88 days. No significant difference between growth performance and liver weight was observed for all diet treatments, except for the diet with 21.7 mg protein/kJ D.P.:D.E., which had significantly lower mean weight gain, feed efficiency and feed conversion.

The digestible protein:digestible energy ratio has become a major factor in fish feed formulation. The Committee on the Nutrient Requirements of Fish and Shrimp of the U.S. National Research has proposed that D.P.:D.E. is a more rational and precise way of conveying crude protein requirements in fish feed. This ratio is important in aquaculture production because a correct ratio will lead to the most efficient conversion of dietary protein into body mass, which maximizes growth and minimizes feed consumption.

Methionine requirements of adult fish

Work by Jonathan Goff and Delbert Gatlin III found that methionine is one of the first limiting essential amino acids in many fish diets, especially those containing higher levels of plant protein sources such as soybean meal, peanut meal and copra meal. With substitution for fishmeal in diets, limiting components such as the sulfur amino acid methionine need to be supplemented.

Many investigators have reported the quantitative methionine requirements of commonly cultured fish species as ranging 18-40 g/kg. This year, J. Niu and co-authors reported the requirement for methionine in golden pompano, *Trachinotus ovatus*, as 10.6 g/kg diet. These results were valuable, but the study was performed on small fish with initial body weights of 12.5 g and final body weights of 72.0 g.

The authors' study focused on determining the methionine requirements for large Florida pompano, *Trachinotus carolinus*, with initial weights of 208 g and final weights of 505 g. These larger fish naturally consumed higher amounts of food. The first results of the investigation suggested that Florida pompano adults perform well as far as growth and feed efficiency with values of methionine below those published by J. Niu.

Perspectives

A closer look at the nutritional requirements and digestive capacity of larger fish is necessary to optimize growth and minimize waste in commercial marine fish farms. These mammoth goals can be more rapidly attained under synergistic collaborations among all stakeholders in the production chain, from suppliers of raw materials and aquafeed manufacturers to commercial farmers and research institutions.

Authors



JORGE A. SUAREZ, PH.D.

University of Miami

Rosenstiel School of Marine and Atmospheric Science

4600 Rickenbacker Causeway

Miami, Florida 33149 USA

jsuarez@rsmas.miami.edu (<mailto:jsuarez@rsmas.miami.edu>).



**CARLOS TUDELA, CHRISTINA BELFRANIN, STEVEN SUTTON, MATTHEW TAYNOR,
RONALD HOENIG, DREW DAVIS, THIAGO RAGGI, DANIEL D. BENETTI, PH.D.**
University of Miami Rosenstiel School of Marine and Atmospheric Science

Copyright © 2016–2019
Global Aquaculture Alliance