



[LEADERSHIP & INNOVATION \(/ADVOCATE/CATEGORY/LEADERSHIP-INNOVATION\)](#)

Philippines farmers apply ASA-IM high-density technology to raise marine fish

Thursday, 1 January 2009

By Levy Loreto L. Manalac , Dr. Michael C. Cremer and Lukas Manomaitis

Low-volume, high-density systems boost productivity, profitability



An important element of the LVHD cage system is to secondarily contain feed within the main cage to minimize waste.

The American Soybean Association – International Marketing (ASA-IM) program Soy-In-Aquaculture project in the Philippines has introduced low-volume, high-density (LVHD) cage culture production methodology for use with high-value marine species. This technology was developed by ASA-IM in China with the aims of maximizing farmers' profits, improving productivity, reducing feed-conversion ratios and limiting environmental degradation.

The system maximizes production through the use of good cage site selection, proper cage positioning, maximum cage volumes and densities, high-quality extruded floating feeds and proper feed management to reduce or prevent disease.

Cage design, orientation

The water quality at LVHD sites can be controlled using a target biomass and specific cage designs and orientation. In marine cages, a final harvest biomass of 50-75 kg/m³ is targeted based on effective water volume enclosed by the cage and surrounding conditions.

The cage design and net mesh size should allow good water exchange inside the cages to maintain dissolved oxygen at a safe level throughout the period of the culture. Cages should be oriented perpendicular to the water current and positioned with at least one cage width between cages in each row. Cage rows should be positioned far enough apart to permit good water exchange.

Feed

One of the most important aspects of a formulated feed-based system is strict feed management using high-quality feeds. Feed is the most expensive part of any production scheme, accounting for 70 to 80 percent of the production cost, and therefore should be used effectively and efficiently.

Soy-In-Aquaculture research has shown that the highest production efficiency is obtained using a floating, extruded feed with a feed enclosure to keep feeds within the cage during feeding. The ASA-IM 90 percent-satiation feeding technique was also developed to help limit wastage of feed.

Introduction of technology

In the Philippines, fish farmers usually used high-volume, low-density cages of 180-1,000 m³ size and fed pelletized sinking feeds ad libitum to produce 10-15 kg/m³ at harvest. They also typically did not change mesh sizes during culture and paid little attention to water exchange in the cage. This led them to believe that a harvest biomass of less than 20 kg/m³ was the maximum they could achieve.

Initially, Philippine fish farmers were very conservative and hesitant about adopting the LVHD technology, and particularly the new feeding techniques using extruded feeds with high-value marine fish species. This conservative attitude was highlighted with two projects in 2007 using pompano, *Trachinotus blochii*, in cage-feeding demonstrations conducted at Alsons Aquaculture Corp. in Davao del Sur, Philippines, and Reprotect Inc. in Negros Oriental, Philippines.

These projects used three 8- or 27-m³ floating cages, respectively, that targeted a final biomass of 50 kg/m³ of fish harvested at the conclusion of the projects using a 43 percent-protein, 12 percent-fat domestically produced, extruded floating aquafeed.

Some of the difficulties seen in adopting the cage production technology were the incorrect application of the 90 percent-satiation feeding technique, improper use of feed enclosures in the cages and particular lack of attention paid to cage mesh size and the water exchange capability of the system, which are critical for good results in a high-density system.



Harvested pompano are packed in chilled ice at Alsons Aquaculture Corp. in Davao Del Sur, Philippines.

Technical support

The ASA-IM program helped show farmers other ways to culture high-value marine fish using smaller cages and higher target densities. It provided close technical support to demonstrate the LVHD cage culture system and continually trained feed managers in the proper way of using the 90 percent-satiation feeding technique throughout the production cycle.

Advice on the proper design and placement of feed enclosures helped prevent floating feed from exiting the cages. This system also taught the farmers the importance of monitoring the relationship of the size of the net mesh to the size of the fish for good water exchange inside the cage.

Perspectives

Despite some challenges encountered in implementing the new technology, the farmers were able to realize the benefits of adopting the new technology compared to their traditional commercial cage culture methods. They had cost savings on both feed consumption and labor using the 90 percent-satiation feeding method without sacrificing fish growth. Using extruded floating feeds helped the farmers better manage their feed use as well as limit environment degradation.

(Editor's Note: This article was originally published in the January/February 2009 print edition of the Global Aquaculture Advocate.)

Authors



LEVY LORETO L. MANALAC

American Soybean Association – International Marketing
37 Bonuan Sabangan, Bonuan Gueset
Dagupan City, Pangasinan
2400 Philippines

levy7172000@yahoo.com (<mailto:levy7172000@yahoo.com>).



DR. MICHAEL C. CREMER

American Soybean Association – International Marketing
St. Louis, Missouri, USA



LUKAS MANOMAITIS

American Soybean Association – International Marketing
Southeast Asia Representative Office
Singapore

Copyright © 2016–2019
Global Aquaculture Alliance