

[ANIMAL HEALTH & WELFARE \(/ADVOCATE/CATEGORY/ANIMAL-HEALTH-WELFARE/\)](/ADVOCATE/CATEGORY/ANIMAL-HEALTH-WELFARE/)

Spermatophore regeneration in black tiger shrimp

Monday, 2 July 2012

By Dr. C.P. Balasubramanian , Dr. P. Ravichandran , D.L. Mohanlal , Dr. S.M. Pillai and Dr. A.G. Ponniah

Spermatophores should be evacuated periodically



Fully melanized and normal spermatophores of *Penaeus monodon*.

Reproduction is a main area where science can contribute to the growth of aquaculture and provide effective solutions for industry problems. Thus, many research projects have focused on the reproductive biology of farmed species. However, most researchers focused on female reproduction – possibly due to the obvious reproductive dysfunction of females under captivity.

In female shrimp, vitellogenesis and oocyte maturation are processes that demand high levels of energy. Conversely, in males, sperm production is believed to consume less energy. Males use up more energy for sex competition and female guarding. Yet male-related reproductive dysfunction and infertility have been increasingly reported in commercial

maturation facilities.

The authors recently carried out a series of studies at the experimental shrimp hatchery facility of the Central Institute of Brackishwater Aquaculture to characterize the reproductive capability of male black tiger shrimp, *Penaeus monodon*, by documenting changes in sperm quality in relation to consecutive regeneration, molting cycle and length of rearing period.

Experimental setup




Fig. 1.: Comparison of sperm counts of *Penaeus monodon* over time. Values with different letters are significantly different.

Thirty broodstock-sized *P. monodon* males were obtained from the broodstock fishery along the coast of Chennai, Tamil Nadu, India. The animals were tagged and reared in the experimental hatchery. Baseline sperm quality characteristics – sperm count, proportion of live sperm and percentage of abnormal sperm – were obtained from freshly extracted

spermatophores.

Spermatophore data and the general condition of the animals were recorded at weekly intervals for 15 animals. The other shrimp remained intact without periodic manual ejaculation. The spermatophores were collected and evaluated on day 30. Further sperm quality was also evaluated at postmolt, intermolt and premolt phases of molting cycle.

Sperm quality variables

While spermatophore weights remained unaffected by consecutive regeneration, sperm count and proportion of viable sperm were better in the subsequent regenerated spermatophores compared to the baseline values (Fig. 1). Further, melanization of spermatophore/male reproductive tracts was not observed. On the contrary, the large proportion of shrimp kept intact without removal of spermatophores developed melanization.

The quality of the melanized spermatophores was extremely poor when compared to the quality of the non-melanized spermatophores (Fig. 2). Histological examination of melanized terminal ampules and testes showed clumps of sperm with melanized debris.

Molt cycle




Fig. 2: Sperm count of *Penaeus monodon* after 30 days under captivity without periodic artificial ejaculation.

Spermatogonia, spermatocytes and spermatids were found in all the phases of the molt cycle. However, only one type of spermatogenic cells was found at any one time. This indicated synchrony in the sperm production. It is also found that the sperm production was a continuous process over the molt cycle (Fig. 3).

Conversely, spermatophore concentration varied during the different molt phases, with significantly higher numbers during the premolt phase. New spermatophores appeared after the exuviations of old exoskeleton.

Perspectives

As suggested by S. Parnes and co-workers in 2006, spermatophores have an inherent “expiration date” and therefore should be evacuated periodically. In the wild, the renovation of spermatophores is carried out by molting or mating. Under captive conditions, mating opportunities and molting frequencies vary.

In the absence of receptive females or failure in molting, the normal renovation of spermatophores does not take place, and they eventually degrade or melanize. Melanization is an end result of prophenoloxidase activation, which is triggered in response to foreign agents such as bacteria, fungi or multicellular parasites, or as a result of tissue damage. Periodic spermatophore renovations by artificial ejaculation are a practical procedure for improving sperm quality in *P. monodon*, as reported in *Litopenaeus* shrimp species.

Sperm quality was found to be highest during the premolt stage, which corresponds to the elevated titre of the molting hormone ecdysteroid. The stimulatory effect of the ecdysteroid on testicular activity of crustaceans has been reported by A. Sagi and co-workers in 1991. The present study indicated that for breeding programs, males at the premolt stage may provide better results than shrimp in other stages.

Editor's Note: Full references are available from the first author.

(Editor's Note: This article was originally published in the July/August 2012 print edition of the Global Aquaculture Advocate.)




Fig. 3: Comparison of sperm count of *Penaeus monodon* at different phases of the molt cycle.

Authors



DR. C.P. BALASUBRAMANIAN

Central Institute of Brackishwater Aquaculture

75 Santhome High Road

R. A. Puram

Chennai 600 028 India

cpbalasubramanian@yahoo.com (<mailto:cpbalasubramanian@yahoo.com>)



DR. P. RAVICHANDRAN

Central Institute of Brackishwater Aquaculture

75 Santhome High Road

R. A. Puram

Chennai 600 028 India



D.L. MOHANLAL
Central Institute of Brackishwater Aquaculture
75 Santhome High Road
R. A. Puram
Chennai 600 028 India



DR. S.M. PILLAI
Central Institute of Brackishwater Aquaculture
75 Santhome High Road
R. A. Puram
Chennai 600 028 India

11/8/2018

Spermatophore regeneration in black tiger shrimp « Global Aquaculture Advocate



DR. A.G. PONNIAH

Central Institute of Brackishwater Aquaculture

75 Santhome High Road

R. A. Puram

Chennai 600 028 India

Copyright © 2016–2018
Global Aquaculture Alliance