

**Alliance**[. \(https://www.aquaculturealliance.org\)](https://www.aquaculturealliance.org)

Health &amp; Welfare

# Testing commercial feeds with tilapia broodstock

Saturday, 1 December 2001

By Ram C. Bhujel, Ph.D.

## Asian Institute of Technology conducts feeding trial in Thailand



Workers collect fry from tilapia reared in hapas.

Nutrient levels in Nile tilapia broodstock feed affect their frequency of spawning, clutch size, egg quality and overall seedstock output. Physical properties of feed – form, type and size of pellets – also have effects on feed intake and thereby reproductive performance.

Specific commercial feeds for tilapia broodstock are generally not available. For other species, only a small number of brood animals are required, so broodstock feed represents a limited market. But for tilapia, for example, 60,000 to 100,000 broodfish are needed to produce 5 to 10 million fry per month. This requires 2 to 4 tons of feed per month for broodstock alone.

Most hatchery operators either prepare their own diets or use feeds made for nursing or fattening the

same or other species, depending upon availability and costs. Farm-made feeds are usually of the sinking type, because extruders are not affordable to fish farmers. These feeds are prone to cause more fouling in hapas, however, so commercial floating feeds are often a better option.

## Testing commercial feeds

Although many types of commercial tilapia feeds are available, it was not known which were best suited for broodstock. Therefore, researchers at the Asian Institute of Technology conducted a feeding trial on a commercial farm in Thailand to compare three locally available commercial feeds for Nile tilapia.

### Feeds

The feeds were different in pellet size, protein content (Table 1), and, of course, price. One feed type was the large, pelleted catfish feed with 25 percent crude protein (CP), the farm was already using.

## Bhujel, Table 1. Composition of commercial feeds, Table 1

	Herbivorous Feed	Catfish Pellets Large	Catfish Pellets Small
Crude Protein Level (%)	15.5	25	30
Fat (%)	4	4	4
Crude Fiber (%)	10	8	8
Moisture (%)	12	12	12
Pellet Size (mm)	5-6	5	3
Feed Price (baht/kg)	8	13.7	15.4

37 baht (Thai currency) = 1 U.S. dollar

Table 1. Composition of commercial feeds used in trial.

Tilapia broodfish are normally grown or bred in greenwater systems, where they receive a considerable portion of nutrients from natural food organisms. Therefore, researchers selected a cheap, low-protein (15.5 percent CP), herbivorous feed. A relatively better quality and easily available small-pellet catfish feed with 30 percent CP was also selected for the trial to compare the reproductive performance of Nile tilapia and the cost effectiveness of the diets.

### Setup

Tests were conducted over 95 days in 12, 120 square meter hapas (24 meters x 5 meters each) placed in two ponds. Nile tilapia (*Oreochromis niloticus*) were stocked at six fish per square meter of hapa space, with an equal mix of male and female animals. Females averaged 92 grams and males averaged 123 grams. The ponds were fertilized weekly at 4 kg nitrogen and 2 kg phosphorus per hectare per day. Animals were fed manually to satiation. The seed were harvested at five-day intervals.

Results as outlined in Table 2, findings included:

- Broodfish ate more small catfish pellets than the other types of feeds.
- Females fed either catfish pellets produced up to 30 percent more eggs and yolk-sac fry than those fed the herbivorous feed.

- The two catfish pellets yielded similar seed output.
- More females spawned from the groups fed large (20 percent) and small (17 percent) catfish pellets than the females fed herbivorous feed.
- Feed type had no effect on the size of eggs and yolksac fry.
- Seed output per gram of protein decreased with the increase in dietary protein.
- Final weights and daily weight gains of both females and males fed small catfish pellets were highest and those fed with the herbivorous diet were lowest.
- Final gonadosomatic index of females (2.57 to 3.22 percent) and males (0.64 to 0.81 percent) was not affected by feed type.
- Production cost per seed was 22 to 23 percent higher for fish fed with the herbivorous diet, as compared to the two catfish pellets.
- Labor accounted for one-third of the total cost of seed production, followed by depreciation of equipment and machines and costs of feed and management.

## Bhujel, Comparative performance of Nile tilapia broodfish, Table 2

	Herbivorous Feed	Catfish Pellets Large	Catfish Pellets Small
Feed Consumption (kg/hapa)	124	135	168
Male Weight Gain (g/fish/day)	1.5	1.9	2.4
Female Weight Gain (g/fish/day)	0.9	1.3	1.8
Clutches/hapa	70	84	82
Seed outputs			
No./m <sup>2</sup> /day	108	138	141
No./g feed	9	10	9
No./g protein	60	43	30
Cost - Seed Production (baht/0.1 million seed)	653	532	536

37 baht (Thai currency) = 1 U.S. dollar

Table 2. Comparative performance of Nile tilapia broodfish fed with three commercial diets.

## Conclusion

The low-quality herbivorous diet produced lower seed output and was less profitable than the two catfish pellets fed to Nile tilapia in a greenwater system. Neither highprotein diets nor higher feeding levels were beneficial in fry production.

At present, most of the hapa-based tilapia hatcheries in Thailand, which produce 1 to 10 million fry per month, use commercial feeds containing 25 to 30 percent crude protein. However, this study showed that 25 percent crude protein diet is nutritionally adequate and cost-effective.

*(Editor's Note: This article was originally published in the December 2001 print edition of the Global Aquaculture Advocate.)*

## Author

---



**RAM C. BHUJEL, PH.D.**

Research Scientist/Post-Doctorate Fellow  
AARM, Asian Institute of Technology, Thailand  
Institute of Aquaculture, Stirling University, United Kingdom

[bhujel@ait.ac.th](mailto:bhujel@ait.ac.th) (<mailto:bhujel@ait.ac.th>)

Copyright © 2016–2020 Global Aquaculture Alliance

All rights reserved.